This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

Application No. 09/701,584. Attorney Docket No. P66095US0

Claim 9 limits claim "II-to embodiments wherein the claimed "support" does not comprise (excludes) "a polyT "spacer."

Claim 20 limits the number of different oligo or polynucleotides to at least 72," support, being found in the present specification at page 4, 3rd complete paragraph.

Claim 21 further limits the number of different oligo- or polynucleotides to "at least 439.".

The number was calculated from the detailed description starting on page 14. This discloses that the glass slide has a size of 76 X 26 mm. It is further disclosed on page 17, line 2, that the distance from one center to the next is 750 µm and 8 dots were obtained for each cDNA (see page 18, line 8).

Claim 23 corresponds to claim 9, with the subject matter preceded by "preferably" being represented in claim 24.

Reconsideration is requested with respect to the rejections of claims 1-10 of record under 35,...

The rejections are based on a combination of Guo et al in view of Chrisey et al. Guo discloses an organic Si-compound bound to a benzene derivative. These two compounds correspond to the bifunctional spacer and the homobifunctional linker of the present invention. The nucleic acid is attached via a polyT-"spacer"; see figure 1 of Guo.

In contrast, thereto, the presently claimed invention avoids the polyT "spacer" and, instead, attaches the oligo- and polynucleotide (i) through a primary amino group attached through an alkane having a length of from 6 to 18 methylene groups or (ii) through a polyether having from 2 to 20 repeating units.

Application No. 09/701,584 Attorney Docket No. P66095US0

As set forth in the papers filed November 9 and December 3, 2002, the method of Guo could not be extended to nucleic acids having a length of 200 to 600 bp. The nucleic acids of Guo are prepared by chemical synthesis. Routine synthesis of nucleic acids cannot reach the length of 200 nucleotides. Therefore, such nucleic acids have to be prepared by an amplification reaction. By using an amplification reaction it is not possible to incorporate the poly T spacer as this spacer would interact with the poly A tail present in normal mRNA.

To summarize it, although Guo discloses a "linker," which corresponds to the bifunctional spacer and bifunctional linker of the presently claimed invention, the chemistry of Guo is not compatible with an amplification procedure for producing the nucleic acids.

Therefore, Guo does not disclose a chemistry suitable for oligonucleotides having a length of 200 to 600 bp, thereby retaining the necessary distance of the oligonucleotide from the surface.

This is done by a polyT "spacer" in Guo and by an alkane or polyether as defined in amended claim

1 of the presently claimed invention.

This deficiency is not overcome by Chrisey et al. This document discloses oligomers having 4 to about 400 bases, typically 20 to about 150 bases and a method for attaching them to a solid support.

It does not disclose any chemistry for attaching oligonucleotides with sufficient distance from the support. Therefore, combining the chemistry of Guo with the disclosure of Chrisey, a person skilled in the art would lack the additional spacer which substitutes the poly/1. "spacer" of Guo.

Application No. 09/701,584
Attorney Docket No. P66095US0

Therefore, one would end up with a different bridge between the solid support and the oligonucleotide. In the definition of figures 1 of Guo, such a combination would lack the "spacer".

A sufficient distance from the solid-support is necessary to give-the-oligonucleotides sufficient flexibility to bind nucleic acids in the solution.

Furthermore, although Chrisey discloses oligonucleotides having 4 to 400 bases, it does not disclose to use nucleotides having a unified length throughout the solid support of 200 to 600 bp as claimed in the presently claimed invention.

Favorable action is requested.

Respectfully submitted,

JACOBSON-HOLMAN PLLC

Ву

William E. Player

Reg. No. 31,409

400 Seventh Street, N.W. The Jenifer Building Washington, D.C. 20004 Tel.: (202) 638-6666

Attorney Docket No. P66095US0

Date: March 3, 2003

WEP/rdt

R:\rthomas\2003\MARCH\P66095USG-Amdr.wpd

ABSTRACT

A support comprising oligo- or polynucleotides covalently linked at their 5'- or 3'-termini to at least one major surface of the support through bifunctional spacers and bifunctional linkers, is characterized in that the oligo- or polynucleotides covalently linked at their 5'- or 3'-termini through bifunctional spacers and bifunctional linkers have a length of from 200 to 600 bp and the oligo- or polynucleotides cambe obtained by a method involving selection of homologous regions of mRNA from a target species and at least one model species; selection of amplification primers allowing the amplification of nucleic acids having a length of from 200 to 600 bp, preferably from 200 to 400 bp, from the homologous regions of both the mRNA from the target species and the mRNA from the at least one model species, the amplification primers optionally having a maximum of 1 mismatch per 6 nucleic acids of the amplification primer; on the at least one major surface of the support, immobilization of the nucleic acids obtained by amplifications of corresponding nucleic acids having, a length of from 200 to 600 bp, for the target species or the at least one model species using the amplification primers.

or work with the